



2-2016

Pull From the Knee: Proper Technique and Application

Brad H. DeWeese

East Tennessee State University

Timothy J. Suchomel

East Stroudsburg University of Pennsylvania

Ambrose J. Serrano

Jarrold D. Burton

Steven K. Scruggs

See next page for additional authors

Follow this and additional works at: http://digitalcommons.sacredheart.edu/pthms_exscifac



Part of the [Exercise Science Commons](#), and the [Sports Sciences Commons](#)

Recommended Citation

DeWeese, B.H., Suchomel, T.J., Serrano, A.J., Burton, J.D., Scruggs, S.K., & Taber, C.B. (2016). Pull From the Knee: Proper Technique and Application. *Strength and Conditioning Journal* 38(1), 79-85. doi: 10.1519/SSC.0000000000000194

This Peer-Reviewed Article is brought to you for free and open access by the Physical Therapy & Human Movement Science at DigitalCommons@SHU. It has been accepted for inclusion in Exercise Science Faculty Publications by an authorized administrator of DigitalCommons@SHU. For more information, please contact ferriby@SacredHeart.edu, lysobeyb@SacredHeart.edu.

Authors

Brad H. DeWeese, Timothy J. Suchomel, Ambrose J. Serrano, Jarrod D. Burton, Steven K. Scruggs, and Christopher B. Taber



The Exercise Technique Column provides detailed explanations of proper exercise technique to optimize performance and safety.

COLUMN EDITOR: Jay Dawes, PhD, CSCS*D, NSCA-CPT*D, FNCSA

Pull From the Knee: Proper Technique and Application

Brad H. DeWeese, EdD,¹ Timothy J. Suchomel, PhD,² Ambrose J. Serrano, MA,³ Jarrod D. Burton, BA,⁴ Steven K. Scruggs, MA,⁵ and Christopher B. Taber, MS¹

¹Center of Excellence for Sport Science and Coach Education, Department of Exercise and Sport Sciences, East Tennessee State University, Johnson City, Tennessee; ²Department of Exercise Science, East Stroudsburg University, East Stroudsburg, Pennsylvania; ³United States Olympic Training Center, Lake Placid, New York; ⁴Wise Athletic Department, University of Virginia's College, Wise, Virginia; and ⁵Department of Physical Education, University of South Carolina, Columbia, South Carolina

ABSTRACT

THE PULL FROM THE KNEE IS A WEIGHTLIFTING MOVEMENT DERIVATIVE THAT CAN BE USED IN THE TEACHING PROGRESSION OF THE CLEAN AND SNATCH EXERCISES. THIS EXERCISE EMPHASIZES POSITIONAL STRENGTH DURING THE TRANSITION PHASE AND THE TRIPLE EXTENSION OF THE HIP, KNEE, AND ANKLE JOINTS THAT IS CHARACTERISTIC OF WEIGHTLIFTING MOVEMENTS.

INTRODUCTION

Much evidence suggests that weightlifting movements are a superior method of training lower-body muscular power compared with power lifting (15), kettlebell (21), and jump training (38). As such, it should come as no surprise that weightlifting movements and their

derivatives are popular exercises that are prescribed by many strength and conditioning practitioners (5,29). Recent literature suggests that weightlifting pulling derivatives that exclude the catch may provide a training stimulus that is comparable (3,4) or superior (31,36,37) to weightlifting derivatives that include the catch. As a result, recent literature has been interested in examining weightlifting pulling derivatives that eliminate the catch phase (6,9,10,28,30–33,35,37). It is clear that researchers and practitioners are interested in examining weightlifting pulling derivatives as a means of training. Given the importance of exercise technique with regard to the training stimulus and injury prevention (17,20), the proper coaching of exercises should not be overlooked.

TYPE OF EXERCISE

The clean pull from the knee (CPK) and snatch pull from the knee (SPK)

are explosive lower-body exercises that can be used to train lower-body muscular power and strength at key positions during the transition and second pull of weightlifting movements. In addition, the CPK and SPK can be used as a part of the teaching progression for the clean and snatch. These exercises can be performed from a static position away of technique blocks or the safety bars of a power rack or with the bar lowered to a hang position at the knee.

MUSCLES INVOLVED

The CPK and SPK involve muscles that have been previously described during similar weightlifting movements (8–11,32,33):

- Static stability in the starting position: Erector spinae group (iliocostalis, longissimus, and spinalis), rectus abdominis, transverse abdominis, external obliques, internal obliques, quadratus lumborum, triceps brachii

(long head), deltoid, subscapularis, latissimus dorsi, brachioradialis, trapezius, splenius capitis, splenius cervicis, infraspinatus, serratus posterior inferior, rhomboid major, rhomboid minor, and the supraspinatus.

- Transition and second pull phases of the CPK and SPK: Upper extremities: trapezius, splenius capitis, splenius cervicis, levator scapulae, rhomboid minor, rhomboid major, serratus posterior superior, posterior deltoid, teres minor, teres major, erector spinae group (iliocostalis, longissimus, and spinalis), rectus abdominis, transverse abdominis, external obliques, and internal obliques. Lower extremities: quadriceps group (rectus femoris, vastus lateralis, vastus medialis, and vastus intermedius), gluteus maximus, hamstrings group (biceps femoris, semimembranosus, and semitendinosus), gastrocnemius, soleus, tibialis posterior, flexor digitorum, peroneus longus, and the peroneus brevis.

BENEFITS OF THE EXERCISE

The SAID principle (Specific Adaptations to Imposed Demands) indicates that a relationship exists between an athlete's training choices and their resultant gains in performance (27). Thus, it is essential for practitioners to prescribe exercises for their athletes that will allow for the greatest transfer to their sport/event. The CPK and SPK are weightlifting pulling derivatives that allow athletes to efficiently transition to the proper peak power position and accelerate the external load using full body triple extension while eliminating the added stress of properly lifting the external load from the floor and catching the bar. Commonalities exist between an athlete's position in the CPK and SPK and sporting movements, such as the shot put, jump shot, tennis serve, max velocity sprinting, and bobsled start (25). Moreover, the static start of these movements may allow for a large transfer of training to benefit athletes who are required to produce high rates of force production and power

from a static starting position (e.g., sprinters in track and field and American football linemen).

Although weightlifting movements typically result in a low injury rate, chronic training with full weightlifting movements involving the catch may result in a greater potential for injury (19,22). A second benefit of the CPK and SPK exercises is that they are less complex with regard to technique and may be used in the teaching progression of the full weightlifting movements. Specifically, these exercises eliminate the catch phase that is characteristic of the full clean and snatch exercises. The previous literature has suggested that the use of weightlifting movement derivatives that eliminate the catch phase may lower the potential for injury (31,34). Furthermore, it has been suggested that weightlifting movements that involve the catch may cause athletes to focus on dropping under the bar to perform the catch, rather than completing the triple extension of the hips, knees, and ankles (36,37). The failure to complete the triple extension movement may then lead to a decrease in the stimulus provided by the exercise.

Because the CPK and SPK eliminate the catch phase of their full movement alternatives, it is possible to overload the transition and triple extension movements to a greater extent than if the catch was performed. By doing this, the athlete may receive a greater training stimulus that may then transfer to an enhanced performance during their respective sport/event. Previous research has indicated that weightlifting pulling derivatives that eliminated the catch phase produced greater kinetic magnitudes and kinematic characteristics (i.e., joint velocities) as compared to weightlifting movements that included the catch phase (3,4,36,37). The primary rationale behind using these movements is that an athlete may be able to train with loads greater than those that

could be used if the catch phase was included.

A final benefit of the CPK and SPK is the possibility of using these movements as potentiating modalities to create greater velocities, likely resulting in greater rates of force development as compared to other weightlifting derivatives. Previous strength-power potentiating complexes have used weightlifting pulling derivatives to enhance subsequent exercise (1,2,26). Owing to the decreased range of motion and ability to overload the triple extension movement, the CPK and SPK may be used as part of a strength-power potentiating complex.

STARTING POSITION- PREPARATION

- The coach or athlete should first set up technique boxes or the safety bars of a squat rack so that the bar is at the appropriate height relative to the athlete's anthropometrics. Specifically, the bar should be positioned directly in front of, but not touching, the knee cap just above the proximal attachment of the patellar tendon.
- After the bar has been positioned properly, the athlete should place their feet approximately hip width apart. The bar should be positioned above the midfoot, and the athlete's toes should be pointed slightly outward to maintain consistent foot positioning with other weightlifting derivatives (8–11,32,33).
- Once the athlete has properly positioned their feet, the athlete should position their hands and grip. The appropriate hand placement will be based on whether the athlete is performing the clean or snatch variation (12). It is recommended that athletes use the "hook grip" (fingers over thumb) for both the CPK and SPK exercises to prevent grip strength being a limiting factor of performance. In addition, because loads in excess of a maximum clean or snatch may be used during the



Figure 1. Starting positions for the clean pull from the knee (left) and snatch pull from the knee (right). Note: The starting position of the athlete may vary slightly based on their anthropometrics.

CPK and SPK, athletes may consider the use of lifting straps.

- After the acquisition of proper hand and grip placement, the athlete should flex forward at the hip while replicating a normal lordotic (i.e., concave) curve in their lumbar spine by isometrically contracting the erector spinae musculature to “raise” the chest. At the same time, the athlete should create and maintain a slight knee bend so that they feel a stretch in their hamstrings. The athlete’s shins should be vertical and perpendicular to the floor, whereas their shoulders should be positioned ahead of the bar.
- The athlete should then be cued to internally rotate their shoulders (glenohumeral joint) and “turn their elbows out” to ensure that a stable arm position exists for the active pulling portion of the CPK and SPK. This will assist in preventing the elbow joint from prematurely bending during the pulling phase of the exercises.
- Athletes should be instructed to “sit on their heels” in the starting position to maximize their ability to produce the greatest possible forces through the platform during the initiation and continuation of the lift. This cue will also allow the athlete to maintain the correct foot pressure during the transition to the peak power position and also will allow greater control and improved bar speed.
- The starting positions of the CPK and SPK are displayed in Figure 1.

TRANSITION TO THE PEAK POWER POSITION

- Before initiating the CPK or SPK from the static starting position, the athlete should create a “tight” torso by creating tension in the muscles of the midsection by inhaling deeply. The athlete should also maintain the lordotic curvature of their lumbar spine to maintain the appropriate



Figure 2. Power positions for the clean pull from the knee (left) and snatch pull from the knee (right). Note: The position of the barbell may vary slightly based on the athlete’s anthropometrics.

hip angle to maximize the force produced into the platform.

- After achieving the set starting position, the athlete should initiate the CPK and SPK by engaging their hamstrings, glutes, and erector spinae muscles to begin to move to the bar vertically.
- During the transition phase of the CPK and SPK, the athlete must transition the bar to the peak power position (9,13,18) to maximize the force and power produced during the later second pull phase of the movement. The transition of the bar from the starting position to the peak power position is accomplished by the athlete by extending

their back while simultaneously moving the hips and knees forward at the same instant and tempo. At this point, the athlete should be moving into a dorsi-flexed position at the ankle joint.

- During the transition phase to the peak power position, the path of the bar should always be moving vertically “up and into” the body.
- The bar should remain as close as possible to the body without touching the thighs until reaching the peak power position (Figures 2 and 3). This will allow the athlete to continue to accelerate the bar without additional frictional influences to slow it down.



Figure 3. Side view of the power position of the clean and snatch pulls from the knee.

SECOND PULL

- The final phase of the CPK and SPK begins as the athlete reaches the power position. As the athlete transitions to the power position, they should use the momentum created during the first pulling action from the knee to build up the intensity into an explosive pull.
- On reaching the power position, the bar should make a “brushing” contact with the thighs before the triple extension movement occurs. The athlete should continue to engage their erector spinae musculature and keep their elbows extended and externally rotated to prevent early bending of the elbows during the pull.
- At this point, the athlete should perform the triple extension movement by explosively extending their hips, knees, and ankles. Simultaneously, the athlete should shrug their shoulders to maximize barbell velocity (Figure 4).
- In addition to the shrug, the athlete should be taught to slightly flex the wrists to keep the barbell close to their body.
- After the pull, the athlete should control the bar’s descent to the mid-thigh position. The athlete can then either lower the bar onto the technique boxes or power rack, or lower the bar to the hang position at the knee in preparation for the next repetition.

COMMON MISTAKES OF THE PULL FROM THE KNEE

- The athlete may begin the movement without their shoulders properly positioned and a rounded back, which may result in an improper transition to the peak power position and may place excess stress on the athlete’s lower back.
- The athlete may not shift to a fully upright position with the shoulders, hips, and midfoot in line before beginning the second pull, causing



Figure 4. The second pull of the clean pull from the knee (left) and snatch pull from the knee (right).

the chest and shoulders to remain ahead of the bar.

- The athlete may not allow the hips and knees to shift forward once the barbell passes the knees (transition phase), likely resulting in a forward pulling motion instead of a vertical pulling motion.
- The athlete may begin the second pull phase too early. Specifically, the athlete will begin the second pull by performing the triple extension motion when the bar is visually too low on the athlete's thigh. This would result in the athlete not properly reaching the necessary power position for maximum force production.
- The athlete may "dip" before beginning the triple extension movement.
- The athlete may push their hips too far forward during the transition and second pull instead of continuing to drive vertically through the midfoot, likely resulting in a looping of the bar away from the athlete's body.
- The athlete may transition their body weight to their forefoot too early, likely resulting in the improper vertical transference of force through the midfoot before the triple extension movement during the second pull.
- The athlete may bend their arms before beginning the second pull, which may prevent maximum transference of generated force to the bar.
- The athlete may not complete the full triple extension movement of the hips, knees, and ankles, ultimately preventing maximum vertical force production.
- The athlete may complete the shrugging motion before the full triple extension movement.
- The athlete may not aggressively shrug at the top of the second pull, preventing maximum bar velocity. Although failing to aggressively shrug at the top of the second pull may not impact lower-body power development, it may impact the transfer of the pulling technique of the CPK and SPK to their catching derivatives.

PRACTICAL APPLICATION

The CPK and SPK weightlifting movements can be implemented in many different training blocks. However, the sets and reps schemes will be determined based on the priority of the training block. For example, the CPK and SPK can be used in a strength-endurance block with a higher repetition range (3 × 10) and lighter-to-moderate loads (7,23,24). Because exercise technique may falter due to fatigue associated with higher repetitions, practitioners implementing the CPK and SPK in a strength-endurance block should consider using cluster sets of 2 or 5 repetitions (14). Loads ranging 80–

100% of the power clean and snatch maximum may be prescribed for the CPK and SPK during a strength-endurance training block. These loading recommendations are based on the decreased displacement of the load that is characteristic of the CPK and SPK compared with the power clean and snatch. The prescription of the CPK and SPK during this training phase may improve the athlete's technique for subsequent training phases with heavier loads and enhance their power-endurance abilities. Practitioners should consider the athlete's abilities before prescribing the CPK and SPK in a strength-endurance block as proper technique may be compromised as a result of fatigue.

Strength and conditioning practitioners may also consider implementing the CPK and SPK into maximal strength and strength-power training blocks using reduced volumes (3 × 5–3 × 3) and increased loads (7,23,24). Because the CPK and SPK exercises do not require the athlete to catch the load after the second pull, practitioners may prescribe loads in excess of 100% of the athlete's maximum clean and snatch. Using a similar weightlifting derivative, Comfort et al. (6) indicated that loads of 120–140% of an athlete's maximum power clean resulted in increases in force production and rate of force development. During a maximal strength or strength-power

phase of training, the CPK and SPK can be used to reinforce technique before transitioning to future training blocks in which the full weightlifting movements may be prescribed.

Finally, the CPK and SPK can be implemented into an explosive speed or maintenance block. When prescribing the CPK and SPK during this phase of training, the primary goal is to enhance power output. This can be accomplished through reduced loads and intensities (3×3 , 3×2 , and 2×2) (7,23,24). Although no previous studies have examined the effect of load on kinetic and kinematic measures during the CPK or SPK, previous research has indicated that peak power output during the midhigh pull occurred at loads of 40–60% of an athlete's maximum power clean (6,16). Owing to the similarities of the CPK and SPK to the midhigh pull, it is likely that similar loads may be prescribed. However, the loads prescribed should be based on the athlete's proficiency and strength. For example, weaker or less technically proficient athletes should focus on improving peak power through lighter loads, whereas heavier loads may be prescribed for a stronger, more technically proficient athlete.

Conflicts of Interest and Source of Funding: The authors report no conflicts of interest and no source of funding.

Brad H. DeWeese is an assistant professor in the Department of Exercise and Sport Sciences at East Tennessee State University.

Timothy J. Suchomel is an assistant professor in the Department of Exercise Science at East Stroudsburg University.

Ambrose J. Serrano is the head strength and conditioning coach at the United States Olympic Training Center at Lake Placid.

Jarrold D. Burton is the strength and conditioning coordinator at University of Virginia's College at Wise.

Steven K. Scruggs is a doctoral student in the Department of Physical Education at the University of South Carolina.

Christopher B. Taber is a doctoral candidate in the Department of Exercise and Sport Sciences at East Tennessee State University.

REFERENCES

- Baker D. The effectiveness of the wave-cycle for in-season training: 20 years of evidence on the in-season maintenance of strength and power in professional athletes. Paper Presented at: 36th National Strength and Conditioning Association Annual Meeting, Las Vegas, NV, July 1, 2013.
- Chiu LZ and Salem GJ. Potentiation of vertical jump performance during a snatch pull exercise session. *J Appl Biomech* 28: 627–635, 2012.
- Comfort P, Allen M, and Graham-Smith P. Comparisons of peak ground reaction force and rate of force development during variations of the power clean. *J Strength Cond Res* 25: 1235–1239, 2011.
- Comfort P, Allen M, and Graham-Smith P. Kinetic comparisons during variations of the power clean. *J Strength Cond Res* 25: 3269–3273, 2011.
- Comfort P, Fletcher C, and McMahon JJ. Determination of optimal loading during the power clean, in collegiate athletes. *J Strength Cond Res* 26: 2970–2974, 2012.
- Comfort P, Udall R, and Jones PA. The effect of loading on kinematic and kinetic variables during the midhigh clean pull. *J Strength Cond Res* 26: 1208–1214, 2012.
- DeWeese BH, Sams M, and Serrano A. Sliding toward Sochi—Part 1: A review of programming tactics used during the 2010-2014 quadrennial. *Natl Strength Cond Assoc Coach* 1: 30–42, 2014.
- DeWeese BH and Scruggs SK. The countermovement shrug. *Strength Cond J* 34: 20–23, 2012.
- DeWeese BH, Serrano AJ, Scruggs SK, and Burton JD. The midhigh pull: Proper application and progressions of a weightlifting movement derivative. *Strength Cond J* 35: 54–58, 2013.
- DeWeese BH, Serrano AJ, Scruggs SK, and Sams ML. The clean pull and snatch pull: Proper technique for weightlifting movement derivatives. *Strength Cond J* 34: 82–86, 2012.
- DeWeese BH, Serrano AJ, Scruggs SK, and Sams ML. The pull to knee—Proper biomechanics for a weightlifting movement derivative. *Strength Cond J* 34: 73–75, 2012.
- Favre M and Peterson MD. Teaching the first pull. *Strength Cond J* 34: 77–81, 2012.
- Haff GG, Stone M, O'Bryant HS, Harman E, Dinan C, Johnson R, and Han KH. Force-time dependent characteristics of dynamic and isometric muscle actions. *J Strength Cond Res* 11: 269–272, 1997.
- Haff GG, Whitley A, McCoy LB, O'Bryant HS, Kilgore JL, Haff EE, Pierce K, and Stone MH. Effects of different set configurations on barbell velocity and displacement during a clean pull. *J Strength Cond Res* 17: 95–103, 2003.
- Hoffman JR, Cooper J, Wendell M, and Kang J. Comparison of Olympic vs. traditional power lifting training programs in football players. *J Strength Cond Res* 18: 129–135, 2004.
- Kawamori N, Rossi SJ, Justice BD, Haff EE, Pistilli EE, O'Bryant HS, Stone MH, and Haff GG. Peak force and rate of force development during isometric and dynamic mid-thigh clean pulls performed at various intensities. *J Strength Cond Res* 20: 483–491, 2006.
- Kraemer WJ, Adams K, Cafarelli E, Dudley GA, Dooly C, Feigenbaum MS, Fleck SJ, Franklin B, Fry AC, Hoffman JR, Newton RU, Potteiger J, Stone MH, Ratamess NA, and Triplett-McBride T; American College of Sports Medicine. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 34: 364–380, 2002.
- Kraska JM, Ramsey MW, Haff GG, Fethke N, Sands WA, Stone ME, and Stone MH. Relationship between strength characteristics and unweighted and weighted vertical jump height. *Int J Sports Physiol Perform* 4: 461–473, 2009.
- Kulund DN, Dewey JB, Brubaker CE, and Roberts JR. Olympic weightlifting injuries. *Phys Sportsmed* 6: 111–119, 1978.
- Lloyd RS, Faigenbaum AD, Stone MH, Oliver JL, Jeffreys I, Moody JA, Brewer C, Pierce KC, McCambridge TM, Howard R, Herrington L, Hainline B, Micheli LJ, Jaques R, Kraemer WJ, McBride MG, Best TM, Chu DA, Alvar BA, and Myer GD. Position statement on youth resistance

- training: The 2014 International Consensus. *Br J Sports Med* 48: 498–505, 2014.
21. Otto WH III, Coburn JW, Brown LE, and Spiering BA. Effects of weightlifting vs. kettlebell training on vertical jump, strength, and body composition. *J Strength Cond Res* 26: 1199–1202, 2012.
 22. Stone MH, Fry AC, Ritchie M, Stoessel-Ross L, and Marsit JL. Injury potential and safety aspects of weightlifting movements. *Strength Cond J* 16: 15–21, 1994.
 23. Stone MH, O'Bryant H, and Garhammer J. A hypothetical model for strength training. *J Sports Med Phys Fitness* 21: 342–351, 1981.
 24. Stone MH, O'Bryant H, Garhammer J, McMillan J, and Rozenek R. A theoretical model of strength training. *Strength Cond J* 4: 36–39, 1982.
 25. Stone MH, Sanborn K, O'Bryant HS, Hartman M, Stone ME, Proulx C, Ward B, and Hruby J. Maximum strength-power-performance relationships in collegiate throwers. *J Strength Cond Res* 17: 739–745, 2003.
 26. Stone MH, Sands WA, Pierce KC, Ramsey MW, and Haff GG. Power and power potentiation among strength-power athletes: Preliminary study. *Int J Sports Physiol Perform* 3: 55–67, 2008.
 27. Stone MH, Stone M, and Sands WA. *Principles and Practice of Resistance Training*. Champaign, IL: Human Kinetics, 2007.
 28. Suchomel TJ, Beckham GK, and Wright GA. Lower body kinetics during the jump shrug: Impact of load. *J Trainology* 2: 19–22, 2013.
 29. Suchomel TJ, Beckham GK, and Wright GA. The impact of load on lower body performance variables during the hang power clean. *Sports Biomech* 13: 87–95, 2014.
 30. Suchomel TJ, Beckham GK, and Wright GA. Effect of various loads on the force-time characteristics of the hang high pull. *J Strength Cond Res* 29: 1295–1301, 2015.
 31. Suchomel TJ, Comfort P, and Stone MH. Weightlifting pulling derivatives: Rationale for implementation and application. *Sports Med* 45: 823–839, 2015.
 32. Suchomel TJ, DeWeese BH, Beckham GK, Serrano AJ, and French SM. The hang high pull: A progressive exercise into weightlifting derivatives. *Strength Cond J* 36: 79–83, 2014.
 33. Suchomel TJ, DeWeese BH, Beckham GK, Serrano AJ, and Sole CJ. The jump shrug: A progressive exercise into weightlifting derivatives. *Strength Cond J* 36: 43–47, 2014.
 34. Suchomel TJ and Sato K. Baseball resistance training: Should power clean variations be incorporated? *J Athl Enhancement* 2, 2013. doi:10.4172/2324-9080.1000112.
 35. Suchomel TJ, Taber CB, and Wright GA. Jump shrug height and landing forces across various loads. *Int J Sports Physiol Perform* 2015. Epub ahead of print.
 36. Suchomel TJ, Wright GA, Kernozek TW, and Kline DE. Kinetic comparison of the power development between power clean variations. *J Strength Cond Res* 28: 350–360, 2014.
 37. Suchomel TJ, Wright GA, and Lottig J. Lower extremity joint velocity comparisons during the hang power clean and jump shrug at various loads. Presented at 32nd International Conference of Biomechanics in Sports, Johnson City, TN, July 2014.
 38. Tricoli V, Lamas L, Carnevale R, and Ugrinowitsch C. Short-term effects on lower-body functional power development: Weightlifting vs. vertical jump training programs. *J Strength Cond Res* 19: 433–437, 2005.

CSCS professionals apply scientific knowledge to train athletes for peak performance.

CSCS is the ONLY accredited strength and conditioning certification.

Being certified by the NSCA means being certified by the best.

The most sought-after strength and conditioning coaches have their CSCS credential.

REACH HIGHER IN YOUR CAREER

NSCA
NATIONAL STRENGTH AND CONDITIONING ASSOCIATION

everyone stronger
NSCA.com